

ON THE K -THEORY OF SUBGROUPS OF VIRTUALLY CONNECTED LIE GROUPS

DANIEL KASPROWSKI

ABSTRACT. We prove that for every finitely generated subgroup G of a virtually connected Lie group which admits a finite-dimensional model for $\underline{E}G$ the assembly map in algebraic K -theory is split injective. We also prove a similar statement for algebraic L -theory, which in particular implies the generalized integral Novikov conjecture for such groups.

1. INTRODUCTION

For every group G and every ring R there is a functor $\mathbb{K}_R: OrG \rightarrow \mathbf{Spectra}$ from the orbit category of G to the category of spectra sending G/H to (a spectrum weakly equivalent to) the K -theory spectrum $\mathbb{K}(R[H])$ for every subgroup $H \leq G$. By K -theory we will always mean non-connective K -theory as constructed by Pedersen and Weibel [PW85]. For any such functor $F: OrG \rightarrow \mathbf{Spectra}$ a G -homology theory $\mathbb{F}$ can be constructed via

$$\mathbb{F}(X) := \mathrm{Map}_G(-, X_+) \wedge_{OrG} F,$$

see Davis and Lück [DL98]. We will denote its homotopy groups by $H_n^G(-, F) := \pi_n \mathbb{F}(X)$. Let $\mathcal{F}$ be a family of subgroups of G . The K -theoretic assembly map for $\mathcal{F}$ is the map

$$\alpha_{\mathcal{F}}: H_n^G(E_{\mathcal{F}}G; \mathbb{K}_R) \rightarrow H_n^G(pt; \mathbb{K}_R) \cong K_n(R[G])$$

induced by the map $E_{\mathcal{F}}G \rightarrow pt$. Here $E_{\mathcal{F}}G$ denote the classifying space for the family $\mathcal{F}$, see Lück [Lüc00]. The assembly map is a helpful tool to related the K -theory of the group ring $R[G]$ to the K -theory of the group rings over $H \in \mathcal{F}$. The assembly map can more generally be defined for any small additive G -category instead of R , see Bartels and Reich [BR07]. In this article all additive categories will be small.

Analogously, for every additive G -category $\mathcal{A}$ with involution and every family of subgroups $\mathcal{F}$ we can define the L -theoretic assembly map

$$\alpha_{\mathcal{F}}: H_n^G(E_{\mathcal{F}}G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) \rightarrow H_n^G(pt; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}).$$

The Farrell–Jones conjecture states that the assembly maps $\alpha_{\mathcal{V}_{cyc}}$ for the family of virtually cyclic subgroups in K - and L -theory are isomorphisms for all additive G -categories $\mathcal{A}$ (with involution) and all $n \in \mathbb{Z}$. It was first formulated in [FJ93]. The Farrell–Jones conjecture has been proved for a large class of groups, for example hyperbolic and CAT(0)-groups, see Bartels and Lück [BL12a, BL12b], Bartels, Lück and Reich [BLR08a, BLR08b] and Wegner [Weg12], virtually solvable groups,

see Wegner [Weg15], and lattices in virtually connected Lie groups, see Bartels, Farrell and Lück [BFL14] and Kammeyer, Lück and Rüping [KLR]. The Farrell–Jones conjecture implies that the assembly maps $\alpha_{\mathcal{F}in}$ for the family of finite subgroups are split injective, see Bartels [Bar03, Theorem 1.3]. The rational split injectivity of the map $\alpha_{\mathcal{F}in}$ in L -theory implies the Novikov conjecture. The integral split injectivity of $\alpha_{\mathcal{F}in}$ is called the *generalized integral Novikov conjecture*, for more details see Section 6. Kasparov proved the Novikov conjecture for all discrete subgroups of virtually connected Lie groups in [Kas88, Theorem 6.9]. More generally, the Novikov conjecture is true for groups which uniformly embed into a Hilbert space, see Skandalis, Tu and Yu [STY02]. This includes all amenable groups and all groups with finite asymptotic dimension. By Carlsson and Goldfarb [CG04, Section 3] and Ji [Ji04, Corollary 3.4] discrete subgroups of virtually connected Lie groups have finite asymptotic dimension giving a second prove that the Novikov conjecture holds for these groups. Here we will in particular show that discrete subgroups of virtually connected Lie groups also satisfy the generalized integral Novikov conjecture.

In [Kas15] the author proved the split injectivity of the assembly map for finitely generated subgroups G of $GL_n(\mathbb{C})$ which have an upper bound on the Hirsch length of the unipotent subgroups. For a definition of the Hirsch length see Section 3. The bound on the Hirsch length exists if and only if G has finite virtual cohomological dimension by Alperin and Shalen [AS81]. Since G is virtually torsion-free, this is the case if and only if there is a finite-dimensional model for $\underline{E}G$ where we consider G with the discrete topology, see Lück [Lüc00, Theorem 3.1]. In this article we want to extend this theorem to subgroups of all virtually connected Lie groups. Note that in the theorem we again consider G with the discrete topology.

Theorem 1.1. *Let G be a finitely generated subgroup of a virtually connected Lie group and assume there exists a finite-dimensional model for $\underline{E}G$. Then the K -theoretic assembly map*

$$H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G])$$

is split injective for every additive G -category $\mathcal{A}$.

A similar version holds for L -theory as well, which in particular implies the generalized integral Novikov conjecture for these groups, see Section 6.

If G is a discrete subgroup of a virtually connected Lie group H and K the maximal compact subgroup of H , then H/K is a finite-dimensional model for $\underline{E}G$, see Lück [Lüc05, Theorem 4.4]. In particular, we get the following corollary.

Corollary 1.2. *Let G be a finitely generated discrete subgroup of a virtually connected Lie group. Then the K -theoretic assembly map*

$$H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G])$$

is split injective for every additive G -category $\mathcal{A}$.

The condition on the existence of a finite-dimensional model for $\underline{E}G$ can be reformulated in the following way.

Proposition 1.3. *A finitely generated subgroup G of a virtually connected Lie group admits a finite-dimensional model for $\underline{E}G$ if and only if there exists $N \in \mathbb{N}$ such that every finitely generated abelian subgroup of G has rank at most N .*

The rank of an abelian group A is defined as $rk(A) := \dim_{\mathbb{Q}}(A \otimes_{\mathbb{Z}} \mathbb{Q})$ or equivalently as the cardinality of maximal linearly independent subset of A . The statement that every finitely generated abelian subgroup of G has rank at most N is equivalent to the statement that every abelian subgroup of G has rank at most N . For a proof of the proposition see [Section 3](#).

In [Section 7](#) we prove that [Theorem 1.1](#) and its L -theoretic analog also hold without the assumption that G is finitely generated.

Acknowledgments: I would like to thank Johannes Ebert for helpful discussions, and Henrik Rüping and the referee for useful comments and suggestions. This work was partially supported by the SFB 878 “Groups, Geometry and Actions” and the Max Planck Society.

2. LIE GROUPS

A Lie group is virtually connected if it has only finitely many connected components. For the rest of this section let H be a virtually connected Lie group with Lie algebra $\mathfrak{h}$ (which we identify with $T_e H$). The Lie group H acts on itself by conjugation

$$c: H \rightarrow \text{Aut}(H), g \mapsto (h \mapsto ghg^{-1}).$$

Taking the derivative yields a map

$$\text{Ad}: H \rightarrow \text{Aut}(\mathfrak{h}), g \mapsto D_e(c(g)).$$

Since $\text{Aut}(\mathfrak{h})$ is a Lie subgroup of $GL(\mathfrak{h})$, Ad gives a representation of H . The kernel of the representation Ad is the centralizer $C_H(H_0)$ of the unit component H_0 of H .

By definition of the centralizer the group $C_H(H_0) \cap H_0$ is abelian and since H is virtually connected the centralizer $C_H(H_0)$ is therefore virtually abelian. For every subgroup G of H we obtain a short exact sequence

$$1 \rightarrow C_H(H_0) \cap G \rightarrow G \rightarrow \text{Ad}(G) \rightarrow 1.$$

with virtually abelian kernel and linear quotient. We will use this sequence to extend the results of [\[Kas15\]](#) to general virtually connected Lie groups. Before we can do so, we first need to prove [Proposition 1.3](#) which will be done in the next chapter.

3. A BOUND ON THE RANK OF ABELIAN SUBGROUPS

In the proof of [Proposition 1.3](#) a bound on the Hirsch-length of the finitely generated nilpotent subgroups is needed. First we review some facts about nilpotent groups to see that this is the same as a bound on the rank of the finitely generated abelian subgroups.

Let G be a group. Define $G_1 := G$ and recursively $G_{n+1} := [G_n, G]$. The series $G = G_1 \geq G_2 \geq \dots$ is called the *lower central series* of G . A group G is *nilpotent* if there exists $c \in \mathbb{N}$ with $G_{c+1} = 1$. The smallest such c is called the *nilpotency class* of G , we denote it by $c(G)$. The *upper central series* $1 = Z_0(G) \leq Z_1(G) \leq \dots$ of G is recursively defined by

$$Z_{i+1}(G) := \{g \in G \mid \forall h \in G : [g, h] \in Z_i(G)\}.$$

If G is nilpotent then $Z_{c(G)}(G) = G$ and the length of the upper and lower central series agree. For any normal subgroup $H \leq G$ the quotient G/H is again nilpotent.

The *Hirsch-length* $h(G)$ of G is

$$h(G) := rk(G_1/G_2) + \dots + rk(G_{c-1}/G_c) + rk(G_c),$$

where $rk(H)$ denotes the rank of an abelian group H , i.e. $rk(H) := \dim_{\mathbb{Q}}(H \otimes_{\mathbb{Z}} \mathbb{Q})$.

Let $n(G)$ denote

$$\max\{rk(A) \mid A \trianglelefteq G \text{ an abelian normal subgroup}\}.$$

Let H be a group and G be a group acting on H . G *acts nilpotently* if there is a series

$$1 = H_0 \leq H_1 \leq \dots \leq H_n = H$$

of G -invariant normal subgroups of H such that the induced action on H_i/H_{i-1} is trivial. In the special case where $H = G$ and the action is by conjugation G acts nilpotently on itself if and only if G is nilpotent.

Proposition 3.1. *Let G be finitely generated nilpotent. Then $h(G) \leq \frac{n(G)(n(G)+1)}{2}$.*

The proposition is proved in Möhres [Möh87, Theorem 2] for torsion-free nilpotent groups instead of finitely generated nilpotent groups. For convenience of the reader we give a proof. For this we need the following well-known statements about nilpotent groups.

Lemma 3.2. *A subgroup of a finitely generated nilpotent group is finitely generated.*

Proof. The statement follows by induction on the nilpotency class. $\square$

Lemma 3.3 ([War76, Theorem 1.3]). *Let G be nilpotent and $N \trianglelefteq G$ a non-trivial normal subgroup. Then $N \cap Z(G)$ is non-trivial, where $Z(G)$ denotes the center of G .*

Lemma 3.4. *Let G be nilpotent and A a maximal abelian normal subgroup. Then $C_G(A) = A$, where $C_G(A)$ is the centralizer of A in G .*

Proof. Since $A \trianglelefteq G$ is normal, so is $C_G(A)$. Suppose $A \neq C_G(A)$. Then $C_G(A)/A$ is a non-trivial normal subgroup of G/A and $H := C_G(A)/A \cap Z(G/A)$ is non-trivial by the previous lemma. Let $C = \langle c \rangle$ be a cyclic subgroup of H . Then $C \trianglelefteq Z(G/A) \trianglelefteq G/A$ and since C lies in the center it is a normal subgroup of G/A . Let $c' \in C_G(A)$ be a pre-image of c , then the pre-image of C is $\langle A, c' \rangle$. This is abelian and normal in G , hence A was not maximal with this property. $\square$

Lemma 3.5. *Let $Tr(n, \mathbb{Z}) \leq GL_n(\mathbb{Z})$ denote the subgroup of unitriangular matrices, i.e. every element of $Tr(n, \mathbb{Z})$ has 1's on the diagonal and 0's below the diagonal. If $G \leq GL_n(\mathbb{Z})$ acts nilpotently on $\mathbb{Z}^n$, then it is unipotent and conjugate to a subgroup of $Tr(n, \mathbb{Z})$.*

Proof. Since $Tr(n, \mathbb{Z})$ is unipotent, it suffices to prove that G is conjugate to a subgroup of it. Let

$$0 = H_0 \trianglelefteq H_1 \trianglelefteq H_2 \trianglelefteq \dots \trianglelefteq H_k = \mathbb{Z}^n$$

be a sequence of G -invariant subspaces and let G act trivially on H_i/H_{i-1} for all $i = 1, \dots, k$. The lemma is obvious for $k = 1$ and we will prove it by induction on k . Let $H' := \{z \in \mathbb{Z}^n \mid \exists l \in \mathbb{Z} : lz \in H_1\}$. Let $z \in H', l \in \mathbb{Z}$ with $lz \in H_1$. For every $g \in G$ we have $lg(z) = g(lz) = lz$ and thus also $g(z) = z$, i.e. G acts

trivially on H' . By construction $\mathbb{Z}^n/H'$ is torsion-free and we obtain a splitting $\mathbb{Z}^n \cong H' \oplus \mathbb{Z}^n/H'$. The sequence

$$0 = H'/H' \trianglelefteq H_2 + H'/H' \trianglelefteq \dots \trianglelefteq H_k + H'/H' = \mathbb{Z}^n/H'$$

consists of G -invariant subspaces and G acts trivially on the quotients. By induction there is a basis of $\mathbb{Z}^n/H$ such that $G \leq GL(\mathbb{Z}^n/H)$ is unitriangular. Using this basis together with a basis of H' yields a basis of $\mathbb{Z}^n$ for which G lies in $Tr(n, \mathbb{Z})$. $\square$

Proof of Proposition 3.1. Let $n := n(G)$ and A be a maximal abelian normal subgroup. Then A again is finitely generated by Lemma 3.2 and $A \cong \mathbb{Z}^n \oplus F$ with F a finite group. G acts by conjugation on A and since $C_G(A) = A$, the induced map $G/A \rightarrow \text{aut}(A)$ is injective. Since F is finite the projection to $\text{aut}(\mathbb{Z}^n) = GL_n(\mathbb{Z})$ has finite kernel. The group G is nilpotent and thus it acts nilpotently on $\mathbb{Z}^n$ (by conjugation). This implies that the image G/A in $GL_n(\mathbb{Z})$ is conjugate to a subgroup of the unitriangular matrices $Tr(n, \mathbb{Z})$. Since $h(Tr(n, \mathbb{Z})) = \frac{n(n-1)}{2}$ we have

$$\begin{aligned} h(G) &\leq h(A) + h(\ker(\text{aut}(A) \rightarrow GL_n(\mathbb{Z}))) + h(Tr(n, \mathbb{Z})) \\ &= n + 0 + \frac{n(n-1)}{2} = \frac{n(n+1)}{2}. \end{aligned}$$

$\square$

A direct corollary of Proposition 3.1 is the following.

Corollary 3.6. *Let G be a group. Then G has a bound on the Hirsch-length of its finitely generated nilpotent subgroups if and only if it has a bound on the rank of its finitely generated abelian subgroups.*

Before we can prove Proposition 1.3 we need the following lemma.

Lemma 3.7. *Let A be a (countable) abelian group with finite rank, then there is a finite-dimensional model for $\underline{E}A$.*

Proof. Let $rk A = n$. Then there exists a subgroup $B \leq A$ isomorphic to $\mathbb{Z}^n$. The quotient $Q := A/B$ has rank 0 and thus is a torsion group. For $n \in \mathbb{N}$ let $F_n \leq Q$ be finite subgroups with $F_n \leq F_{n+1}$ and $Q = \bigcup_{n \in \mathbb{N}} F_n$. Define a Q -CW complex X by taking $\coprod_{n \in \mathbb{N}} Q/F_n$ as the zero skeleton and for every $n \in \mathbb{N}$ adding a 1-cell with stabilizer F_n between the 0-cells Q/F_n and Q/F_{n+1} . This defines a 1-dimensional model X for $\underline{E}Q$. Let $p: A \rightarrow Q$ be the quotient map. For every finite subgroup $F \leq Q$ the preimage $p^{-1}(F)$ is finitely generated abelian of rank n and thus has $\mathbb{R}^n$ as an n -dimensional model for $\underline{E}p^{-1}(F)$. Therefore, the proof of Lück [Lüc00, Theorem 3.1] shows that A has a model for $\underline{E}A$ of dimension $n + 1$. $\square$

Let G be a subgroup of $GL_n(\mathbb{C})$ and assume there exists $N \in \mathbb{N}$ such that the rank of every finitely generated unipotent subgroup of G is at most N . Then by Alperin and Shalen [AS81] the virtual cohomological dimension of G is bounded and therefore admits a finite-dimensional model for $\underline{E}G$ by [Lüc00, Theorem 6.4]. Using this we now can prove Proposition 1.3.

Proof of Proposition 1.3. Let G be a subgroup of a virtually connected Lie group H such that there exists a finite dimensional model X for $\underline{E}G$. Then in particular X is a model for $\underline{E}A$ for every abelian subgroup $A \leq G$ and $rk A \leq \dim X$.

For the other direction let G be a finitely generated subgroup of a virtually connected Lie group H such that there exists a bound on the rank of the finitely generated abelian subgroups of G . Then by [Corollary 3.6](#) G has also a bound on the Hirsch-length of its finitely generated nilpotent subgroups. Let $G_0 := G \cap H_0$ and consider the extension

$$1 \rightarrow C_H(H_0) \cap G_0 \rightarrow G_0 \rightarrow \text{Ad}(G_0) \rightarrow 1$$

from [Section 2](#). Since $C_H(H_0) \cap G_0$ is contained in the center of G_0 also $\text{Ad}(G_0)$ has a bound on the the Hirsch-length of its finitely generated nilpotent subgroups and thus on the finitely generated unipotent subgroups. By the above it admits a finite dimensional model for $\underline{\text{EAd}}(G_0)$. And since also $K := C_H(H_0) \cap G_0$ has finite rank, there is a finite dimensional model for $\underline{\text{EK}}$ by [Lemma 3.7](#). Consider the extensions

$$1 \rightarrow K \rightarrow G_0 \rightarrow \text{Ad}(G_0) \rightarrow 1$$

and

$$1 \rightarrow G_0 \rightarrow G \rightarrow F \rightarrow 1$$

with F finite. The group G_0 is finitely generated since finite index subgroups of finitely generated groups are again finitely generated. Thus $\text{Ad}(G_0)$ is virtually torsion-free by Selberg's Lemma and we can use [\[Lüc00, Theorem 3.1\]](#) to obtain a finite dimensional model for $\underline{\text{EG}}$ from these sequences. $\square$

Remark 3.8. Using the results of the author from [\[Kas15\]](#) the short exact sequence

$$1 \rightarrow C_H(H_0) \cap G \rightarrow G \rightarrow \text{Ad}(G) \rightarrow 1$$

implies that G has fqFDC, which also is defined in [\[Kas15\]](#). In particular, if G has a bound on the order of the finite subgroups, then the main result of [\[Kas15\]](#) directly implies the split injectivity of the K -theoretic assembly map and a similar result in L -theory. Since we do not know if this always holds, we use a different approach using inheritance properties, see [Sections 4 and 5](#).

4. INHERITANCE PROPERTIES

To use the short exact sequence from [Section 2](#) we want to show the following inheritance property.

Proposition 4.1. *Assume there is a short exact sequence of groups*

$$1 \rightarrow J \rightarrow G \xrightarrow{\phi} Q \rightarrow 1$$

such that for every virtually cyclic subgroup $V \leq Q$ the pre-image $\phi^{-1}(V)$ satisfies the Farrell–Jones conjecture. Furthermore, assume that the assembly map

$$H_n^G(\underline{\text{EQ}}; \mathbb{K}_{\mathcal{B}}) \rightarrow K_n(\mathcal{B}[Q])$$

is split injective for every $n \in \mathbb{Z}$ and every additive Q -category $\mathcal{B}$. Then the K -theoretic assembly map

$$H_n^G(\underline{\text{EG}}; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G])$$

is split injective for every $n \in \mathbb{Z}$ and every additive G -category $\mathcal{A}$.

Proof. Let $\mathcal{A}$ be an additive G -category. That $\phi^{-1}(V)$ satisfies the Farrell–Jones conjecture for every virtually cyclic subgroup $V \leq Q$ implies that the natural map $H_n^G(E_{\mathcal{V}cyc}G; \mathbb{K}_{\mathcal{A}}) \rightarrow H_n^G(E_{\phi^*\mathcal{V}cyc}G; \mathbb{K}_{\mathcal{A}})$ is an isomorphism by Bartels and Lück [BL06, Lemma 2.2], where $\phi^*\mathcal{V}cyc := \{K \leq G \mid \phi(K) \in \mathcal{V}cyc\}$. Here we used that the projection $E_{\mathcal{V}cyc}G \times E_{\phi^*\mathcal{V}cyc}G \rightarrow E_{\phi^*\mathcal{V}cyc}G$ is a model for the natural map $E_{\mathcal{V}cyc}G \rightarrow E_{\phi^*\mathcal{V}cyc}G$. Furthermore, the natural map $H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow H_n^G(E_{\mathcal{V}cyc}G; \mathbb{K}_{\mathcal{A}})$ is split injective by Bartels [Bar03]. Now the commutative diagram

$$\begin{array}{ccc} H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) & \longrightarrow & H_n^G(E_{\phi^*\mathcal{F}in}G; \mathbb{K}_{\mathcal{A}}) \\ \downarrow & & \downarrow \\ H_n^G(E_{\mathcal{V}cyc}G; \mathbb{K}_{\mathcal{A}}) & \xrightarrow{\cong} & H_n^G(E_{\phi^*\mathcal{V}cyc}G; \mathbb{K}_{\mathcal{A}}) \end{array}$$

implies that the map $H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow H_n^G(E_{\phi^*\mathcal{F}in}G; \mathbb{K}_{\mathcal{A}})$ is split injective, where $\phi^*\mathcal{F}in := \{K \leq G \mid \phi(K) \in \mathcal{F}in\}$. By Bartels and Reich [BR07, Corollary 4.3] the split injectivity for Q implies that the assembly map $H_n^G(E_{\phi^*\mathcal{F}in}G; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G])$ is split injective. Combining these results yields the proposition. $\square$

To apply the above proposition for the short exact sequence from the previous section we need the following.

Lemma 4.2. *The class of virtually solvable groups is closed under group extensions.*

The idea of the proof is taken from math.stackexchange.com, see [dC].

Proof. Let

$$1 \rightarrow N \rightarrow G \xrightarrow{p} Q \rightarrow 1$$

be a short exact sequence and let N, Q be virtually solvable. Let $Q' \leq Q$ be a solvable subgroup with $[Q : Q'] < \infty$, then $[G : p^{-1}(Q')] < \infty$. Thus we can assume that Q is solvable. We will first consider the case that N is finite. Since N is normal in G , G acts on N by conjugation, which induces a map $c: G \rightarrow \text{Aut}(N)$. The centralizer $C_G(N)$ of N in the G is the kernel of c . Since the class of solvable groups is closed under extension and $C_G(N) \cap N$ is abelian the exact sequence

$$1 \rightarrow C_G(N) \cap N \rightarrow C_G(N) \rightarrow p(C_G(N)) \rightarrow 1$$

shows that $C_G(N)$ is solvable. The group N is finite, thus $C_G(N)$ has finite index in G .

Now let N be any virtually solvable group. And let $\mathcal{S}$ be the set of all normal, solvable, finite-index subgroups of N ordered by inclusion. This is not empty and we can choose K to be a maximal element of $\mathcal{S}$. For every $g \in G$ also $gKg^{-1}K$ is a solvable, normal, finite-index subgroup of N . Since K was maximal it therefore has to be normal in G . From the short exact sequence

$$1 \rightarrow N/K \rightarrow G/K \rightarrow Q \rightarrow 1$$

it follows from the first case that G/K is virtually solvable. Since K is solvable the sequence

$$1 \rightarrow K \rightarrow G \rightarrow G/K \rightarrow 1$$

implies that G is virtually solvable. $\square$

5. PROOF OF THEOREM 1.1

For this section let H be a virtually connected Lie group and $G \leq H$ a finitely generated subgroup such that there exists a finite dimensional model for $\underline{E}G$. The proof of Theorem 1.1 follows easily from the statements of the previous section.

Proof of Theorem 1.1. Let $\Gamma := \text{Ad}(G)$ be the image of G under $\text{Ad}: H \rightarrow GL(\mathfrak{h})$. Since $C_H(H_0) \cap G \cap H_0$ is contained in the center of G , the pre-image of any unipotent subgroup U of $\text{Ad}(G \cap H_0)$ is a nilpotent subgroup of $G \cap H_0$. By Corollary 3.6 and Proposition 1.3 there is a bound on the Hirsch-length of the nilpotent subgroups of $G \cap H_0$ and in particular there is a bound on the Hirsch-length of U . Since $G \cap H_0$ has finite index in G this implies that there also is a bound on the Hirsch-length of the unipotent subgroups of Γ . Now we can apply

[Kas15, Corollary 3]. *Let F be a field of characteristic zero, Γ a finitely generated subgroup of $GL_n(F)$ with a global upper bound on the Hirsch rank of its unipotent subgroups. Then the K -theoretic assembly map*

$$H_*^\Gamma(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow H_*^\Gamma(pt; \mathbb{K}_{\mathcal{A}}) \cong K_*(\mathcal{A}[\Gamma])$$

is split injective for every additive Γ -category $\mathcal{A}$.

Note that [Kas15, Corollary 3] is stated only for rings instead of additive Γ -categories, but by [Kas15, Theorem 8.1] it is true for any additive Γ -category.

Furthermore, by Wegner [Weg15] every virtually solvable group satisfies the Farrell–Jones conjecture. Using this and Lemma 4.2 we see that the sequence

$$1 \rightarrow C_H(H_0) \cap G \rightarrow G \rightarrow \text{Ad}(G) \rightarrow 1$$

satisfies the conditions of Proposition 4.1. Therefore, the assembly map

$$H_*^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow K_*(\mathcal{A}[G])$$

is split injective for every additive G -category $\mathcal{A}$. □

6. L-THEORY

Most of the statements from the previous sections also hold for L -theory. For the rest of the section let G be a finitely generated subgroup of a virtually-connected Lie group H with a finite dimensional model for $\underline{E}G$ and let Q be the image of G under $\text{Ad}: H \rightarrow GL(\mathfrak{h})$. Furthermore, let ϕ denote $\text{Ad}|_G$ and let $\mathcal{A}$ be an additive G -category with involution. As above we obtain the commutative diagram

$$\begin{array}{ccc} H_n^G(\underline{E}G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) & \longrightarrow & H_n^G(E_{\phi^* \text{Fin}} G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) \\ \downarrow & & \downarrow \\ H_n^G(E_{\mathcal{V} \text{cyc}} G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) & \xrightarrow{\cong} & H_n^G(E_{\phi^* \mathcal{V} \text{cyc}} G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) \end{array}$$

and the lower horizontal map is still an isomorphism by Bartels and Lück [BL06, Lemma 2.2] and Wegner [Weg15]. But for the vertical map on the left to be injective we need that for every virtually cyclic subgroup $V \subseteq G$ there is an $i_0 \in \mathbb{N}$ such that for every $i \geq i_0$ we have $K_{-i}(\mathcal{A}[V]) = 0$, see Bartels [Bar03]. Then it remains to show that

$$H_n^G(E_{\phi^* \text{Fin}} G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) \rightarrow L_n^{\langle -\infty \rangle}(\mathcal{A}[G])$$

is split injective. By Bartels and Reich [BR07, Proposition 4.2 and Corollary 4.3] this follows if

$$H_n^Q(\underline{E}G; \mathbb{L}_{\text{ind}_\phi \mathcal{A}}^{\langle -\infty \rangle}) \rightarrow L_n^{\langle -\infty \rangle}((\text{ind}_\phi \mathcal{A})[Q])$$

is split injective. See [BR07] for the definition of $\text{ind}_\phi \mathcal{A}$. To apply [Kas15, Theorem 9.1] as above, we need the further assumption that for every finite subgroup $H \leq Q$ there is an $i_0 \in \mathbb{N}$ such that for every $i \geq i_0$ we have

$$0 = K_{-i}((\text{ind}_\phi \mathcal{A})[H]) \cong K_{-i}(\mathcal{A}[\phi^{-1}(H)]).$$

Since $\phi^{-1}(H)$ is virtually abelian we obtain the following version of the main theorem for L -theory.

Theorem 6.1. *Let G be a finitely generated subgroup of a virtually connected Lie group and assume there exists an $N \in \mathbb{N}$ such that every finitely generated abelian subgroup of G has at most rank N . Let $\mathcal{A}$ be an additive G -category with involution. Assume further that for every virtually abelian subgroup H of G there is an $i_0 \in \mathbb{N}$ such that for every $i \geq i_0$ we have $K_{-i}(\mathcal{A}[H]) = 0$, then the L -theoretic assembly map*

$$H_n^G(\underline{E}G; \mathbb{L}_{\mathcal{A}}^{\langle -\infty \rangle}) \rightarrow L_n^{\langle -\infty \rangle}(\mathcal{A}([G]))$$

is split injective.

For torsion-free groups G the *integral Novikov conjecture* states that the assembly map

$$H_n^G(EG; \mathbb{L}_{\mathbb{Z}}^{\langle -\infty \rangle}) \rightarrow L_n^{\langle -\infty \rangle}(\mathbb{Z}[G])$$

is injective. It is known that the integral Novikov conjecture is false for groups containing torsion. Following Ji [Ji07] we say that G satisfies the *generalized integral Novikov conjecture* if the assembly maps

$$H_n^G(\underline{E}G; \mathbb{L}_{\mathbb{Z}}^{\langle -\infty \rangle}) \rightarrow L_n^{\langle -\infty \rangle}(\mathbb{Z}[G]), \quad H_n^G(\underline{E}G; \mathbb{K}_{\mathbb{Z}}) \rightarrow K_n(\mathbb{Z}[G])$$

are injective. By Lück and Reich [LR05, Propostion 2.20] the relative rational assembly map

$$H_n^G(EG; \mathbb{L}_{\mathbb{Z}}^{\langle -\infty \rangle}) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow H_n^G(\underline{E}G; \mathbb{L}_{\mathbb{Z}}^{\langle -\infty \rangle}) \otimes_{\mathbb{Z}} \mathbb{Q}$$

is injective. Observe that, since the $\mathbb{Z}/2$ -Tate cohomology groups vanish rationally, there is no difference between the various decorations in L -theory because of the Rothenberg sequence. Therefore, by [LR05, Proposition 1.53] the injectivity of the rational assembly map

$$H_n^G(EG; \mathbb{L}_{\mathbb{Z}}^{\langle -\infty \rangle}) \otimes_{\mathbb{Z}} \mathbb{Q} \rightarrow L_n^{\langle -\infty \rangle}(\mathbb{Z}[G]) \otimes_{\mathbb{Z}} \mathbb{Q}$$

implies the Novikov conjecture about the homotopy invariance of higher signatures. In particular, the generalized integral Novikov conjecture implies the (classical) Novikov conjecture.

We will show that $K_{-n}(\mathbb{Z}[G]) = 0$ for $n > 1$ and any virtually abelian group A . Therefore, Theorem 6.1 implies the generalized integral Novikov conjecture for the groups G appearing in the theorem, i.e. we get the following corollary.

Corollary 6.2. *Let G be a finitely generated subgroup of a virtually connected Lie group and assume there exists an $N \in \mathbb{N}$ such that every finitely generated abelian subgroup of G has at most rank N . Then G satisfies the generalized integral Novikov conjecture.*

By Farrell and Jones [FJ95, Theorem 2.1] for every virtually cyclic group V and $n > 1$

$$K_{-n}(\mathbb{Z}[V]) = 0.$$

Let G be a group and let X be a finite G -CW-complex with virtually cyclic stabilizers. By induction on the dimension of X we prove that

$$H_{-n}^G(X; \mathbb{K}_{\mathbb{Z}}) = 0$$

for every $n > 1$. For $\dim X = 0$ we have

$$H_{-n}^G(X; \mathbb{K}_{\mathbb{Z}}) \cong \bigoplus_{x \in X} \mathbb{K}_{-n}(\mathbb{Z}[G_x]) = 0,$$

where the stabilizers G_x are virtually cyclic by assumption. Assume the above holds for m and let $\dim X = m + 1$. Then we have the long exact sequence

$$0 = H_{-n}^G(X^{(m)}; \mathbb{K}_{\mathbb{Z}}) \rightarrow H_{-n}^G(X; \mathbb{K}_{\mathbb{Z}}) \rightarrow H_{-n}^G(X, X^{(m)}; \mathbb{K}_{\mathbb{Z}})$$

and

$$H_{-n}^G(X, X^{(m)}; \mathbb{K}_{\mathbb{Z}}) \cong \bigoplus_{c \in C_m} \mathbb{K}_{-n-m-1}(\mathbb{Z}[G_c]) = 0,$$

where C_m denotes the set of m -cells of X and G_c the (virtually-cyclic) stabilizer of the cell c . Since every virtually abelian group A satisfies the Farrell–Jones conjecture we have

$$K_{-n}(\mathbb{Z}[A]) \cong H_{-n}^A(X; \mathbb{K}_{\mathbb{Z}}) \cong \operatorname{colim}_K H_{-n}^A(AK; \mathbb{K}_{\mathbb{Z}}) = 0,$$

where X is a A -CW-complex model for $E_{\mathcal{V}cyc}A$ and the colimit is taken over all finite subcomplexes $K \subseteq X$.

7. INHERITANCE UNDER COLIMITS

In this section we want to show that [Theorem 1.1](#) and [Theorem 6.1](#) hold without the assumption that G is finitely generated.

By Bartels, Echterhoff and Lück [BEL08, Lemma 2.4, Lemma 6.2] for every system G_α of finitely generated subgroups of G such that $\operatorname{colim}_\alpha G_\alpha \cong G$ the assembly map

$$H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G])$$

is the colimit of the assembly maps

$$H_n^{G_\alpha}(\underline{E}G_\alpha; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G_\alpha]),$$

for any additive G -category $\mathcal{A}$. The same statement holds in L -theory for any additive G -category with involution. Note that the statement in [BEL08] is formulated for rings with G -action instead of additive G -categories, but the statement for G -categories holds in the same way. Furthermore, a finite-dimensional model for $\underline{E}G$ gives a finite-dimensional model for $\underline{E}G_\alpha$ by restricting the action to G_α . So taking the colimit over all finitely generated subgroups proves that injectivity holds without the assumption that G is finitely generated. For the construction of a splitting we need to see that the splittings for the finitely generated subgroups are natural with respect to the structure maps of the colimit. In the proof of [Theorem 1.1](#) and [Theorem 6.1](#) the assumption that G is finitely generated is only needed to apply [Kas15, Corollary 3] respectively its L-theoretic analog. So it suffices to see that the splittings constructed in [Kas15] are natural with respect to the structure maps of the colimit.

We will use the definitions of controlled categories and bounded mapping spaces from [Kas15, Section 5 and 7]. In the following let X denote a finite dimensional simplicial model for $\underline{E}G$. By Bartels, Farrell, Jones and Reich [BFJR04, Section 6] the assembly map

$$H_n^G(\underline{E}G; \mathbb{K}_{\mathcal{A}}) \rightarrow K_n(\mathcal{A}[G])$$

can be identified with the map

$$\operatorname{colim}_{K \subseteq X \text{ fin.}} \pi_{n+1}(\mathbb{K}\mathcal{A}_G(GK)^\infty)^G \rightarrow \operatorname{colim}_{K \subseteq X \text{ fin.}} \pi_n(\mathbb{K}\mathcal{A}_G(GK)_0)^G.$$

Now consider the diagram

$$\begin{array}{ccc} \operatorname{colim}_{K \subseteq X \text{ fin.}} \pi_{n+1}(\mathbb{K}\mathcal{A}_G(GK)^\infty)^G & \xrightarrow{\quad} & \operatorname{colim}_{K \subseteq X \text{ fin.}} \pi_n(\mathbb{K}\mathcal{A}_G(GK)_0)^G \\ \downarrow f & & \downarrow j \\ \operatorname{colim}_{K \subseteq X \text{ fin.}} \pi_{n+1} \operatorname{Map}_G^{bd}(X, \mathbb{K}\mathcal{A}_G(GK)^\infty) & \xrightarrow{h} & \operatorname{colim}_{K \subseteq X \text{ fin.}} \pi_n \operatorname{Map}_G^{bd}(X, \mathbb{K}\mathcal{A}_G(GK)_0). \end{array}$$

By [Kas15, Remark 7.7] the map f is an isomorphism and the map h is an isomorphism in the situation of [Kas15, Corollary 3].

Let $\Gamma \rightarrow \Lambda$ be an injective group homomorphism. For every Λ -set J and every subcomplex $K \subseteq X$ we can define a map

$$\left(\prod_J^{bd} \mathcal{A}_\Gamma(\Gamma K)^\infty \right)^\Gamma \rightarrow \left(\prod_J^{bd} \mathcal{A}_\Lambda(\Lambda K)^\infty \right)^\Lambda$$

as follows. A controlled modul $(M_j) \in \left(\prod_J^{bd} \mathcal{A}_\Gamma(\Gamma K) \right)^\Gamma$ is sent to $(M'_j)_j$ with $(M'_j)_{h',x,t} := \bigoplus_{[h] \in \Lambda/\Gamma} (M_{h^{-1}j})_{h^{-1}h',h^{-1}x,t}$ and analogously on morphisms. This map is well defined since (M_j) is Γ -invariant. The above maps induce a map

$$\operatorname{Map}_\Gamma^{bd}(X, \mathbb{K}\mathcal{A}_\Gamma(\Gamma K)) \rightarrow \operatorname{Map}_\Lambda^{bd}(X, \mathbb{K}\mathcal{A}_\Lambda(\Lambda K))$$

for every finite subcomplex $K \subseteq X$ and in the special case where $J = \{pt\}$ we obtain a map

$$(\mathbb{K}\mathcal{A}_\Gamma(\Gamma K)^\infty)^\Gamma \rightarrow (\mathbb{K}\mathcal{A}_\Lambda(\Lambda K)^\infty)^\Lambda.$$

The same maps can be constructed with $\mathcal{A}_\Gamma(\Gamma K)^\infty$ and $\mathcal{A}_\Lambda(\Lambda K)^\infty$ replaced by $\mathcal{A}_\Gamma(\Gamma K)_0$ and $\mathcal{A}_\Lambda(\Lambda K)_0$ respectively. So they induce maps from the above diagram for Γ to the same diagram for Λ . We will omit the technical proofs that the maps of the diagram are natural with respect to these maps and that under the identification with the assembly map they correspond to the structure maps of the colimit from [BFJR04]. This shows that the splitting $f^{-1} \circ h^{-1} \circ j$ is natural with respect to the structure maps of the colimit.

Now let us consider the L -theoretic version. For [Kas15, Remark 7.7] it was used that the category $(\prod_{j \in J} \mathbb{K}\mathcal{A}_G(GK)^\infty)^G \simeq \prod_{[j] \in G \backslash J} \mathbb{K}\mathcal{A}_G^{G_j}(GK)^\infty$ is weakly equivalent to $(\mathbb{K} \prod_{j \in J} \mathcal{A}_G(GK)^\infty)^G \simeq \mathbb{K} \prod_{[j] \in G \backslash J} \mathcal{A}_G^{G_j}(GK)^\infty$ for every G -set J with finite stabilizers and every finite subcomplex $K \subseteq X$, where G_j is the stabilizer of $j \in J$. Let $H \leq G$ be finite, then

$$K_n(\mathcal{A}_G^{G_j}(G/H)^\infty) \cong \prod_{G_j \backslash G/H} K_n(\mathcal{A}_G^{G_j}(G_j/(G_j \cap H))^\infty) \cong \prod_{G_j \backslash G/H} K_{n-1}(\mathcal{A}[G_j \cap H]).$$

If for each finite subgroup $H \leq G$ there exists $N \in \mathbb{N}$ such that for each $n \geq N$ the groups $K_{-n}\mathcal{A}[H]$ vanish, then by induction on the cells this implies that for

every finite subcomplex $K \subseteq X$ there exists $N \in \mathbb{N}$ such that for $n \geq N$ the groups $K_{-n}(\mathcal{A}_G^{G_j}(GK)^\infty)$ vanish. Therefore, under this assumption L -theory commutes with the above product and we get that the map

$$\phi: \operatorname{Map}_G^{bd}(X, \mathbb{L}\mathcal{A}_G(GK)^\infty) \rightarrow \operatorname{Map}_G(X, \mathbb{L}\mathcal{A}_G(GK)^\infty)$$

is an isomorphism and under the above assumption also

$$\psi: (\mathbb{L}\mathcal{A}_G(GK)^\infty)^G \rightarrow \operatorname{Map}_G(X, \mathbb{L}\mathcal{A}_G(GK)^\infty)$$

is an isomorphism, see [Kas15, Section 9]. Since ψ factors over ϕ the map

$$(\mathbb{L}\mathcal{A}_G(GK)^\infty)^G \rightarrow \operatorname{Map}_G^{bd}(X, \mathbb{L}\mathcal{A}_G(GK)^\infty)$$

is an isomorphism as well. Therefore, we obtain the naturality of the splitting as in the case for K -theory.

REFERENCES

- [AS81] Roger C. Alperin and Peter B. Shalen, *Linear groups of finite cohomological dimension*, Bull. Amer. Math. Soc. (N.S.) **4** (1981), no. 3, 339–341. MR 609046 (82c:20087)
- [Bar03] Arthur C. Bartels, *On the domain of the assembly map in algebraic K-theory*, Algebr. Geom. Topol. **3** (2003), 1037–1050. MR 2012963 (2004h:19002)
- [BEL08] Arthur Bartels, Siegfried Echterhoff, and Wolfgang Lück, *Inheritance of isomorphism conjectures under colimits*, K-theory and noncommutative geometry, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2008, pp. 41–70. MR 2513332 (2012a:55005)
- [BFJR04] Arthur Bartels, Tom Farrell, Lowell Jones, and Holger Reich, *On the isomorphism conjecture in algebraic K-theory*, Topology **43** (2004), no. 1, 157–213. MR 2030590 (2004m:19004)
- [BFL14] A. Bartels, F. T. Farrell, and W. Lück, *The Farrell-Jones conjecture for cocompact lattices in virtually connected Lie groups*, J. Amer. Math. Soc. **27** (2014), no. 2, 339–388. MR 3164984
- [BL06] Arthur Bartels and Wolfgang Lück, *Isomorphism conjecture for homotopy K-theory and groups acting on trees*, J. Pure Appl. Algebra **205** (2006), no. 3, 660–696. MR 2210223 (2007e:19005)
- [BL12a] ———, *The Borel conjecture for hyperbolic and CAT(0)-groups*, Ann. of Math. (2) **175** (2012), no. 2, 631–689. MR 2993750
- [BL12b] ———, *Geodesic flow for CAT(0)-groups*, Geom. Topol. **16** (2012), no. 3, 1345–1391. MR 2967054
- [BLR08a] Arthur Bartels, Wolfgang Lück, and Holger Reich, *Equivariant covers for hyperbolic groups*, Geom. Topol. **12** (2008), no. 3, 1799–1882. MR 2421141 (2009d:20102)
- [BLR08b] ———, *The K-theoretic Farrell-Jones conjecture for hyperbolic groups*, Invent. Math. **172** (2008), no. 1, 29–70. MR 2385666 (2009c:19002)
- [BR07] Arthur Bartels and Holger Reich, *Coefficients for the Farrell-Jones conjecture*, Adv. Math. **209** (2007), no. 1, 337–362. MR 2294225 (2008a:19002)
- [CG04] Gunnar Carlsson and Boris Goldfarb, *On homological coherence of discrete groups*, J. Algebra **276** (2004), no. 2, 502–514. MR 2058455 (2005a:20078)
- [dC] Yves de Cornulier, *If N and G/N are virtually solvable, then G is virtually solvable?*, (version: 2013-07-05).
- [DL98] James F. Davis and Wolfgang Lück, *Spaces over a category and assembly maps in isomorphism conjectures in K- and L-theory*, K-Theory **15** (1998), no. 3, 201–252. MR 1659969 (99m:55004)
- [FJ93] F. T. Farrell and L. E. Jones, *Isomorphism conjectures in algebraic K-theory*, J. Amer. Math. Soc. **6** (1993), no. 2, 249–297. MR 1179537 (93h:57032)
- [FJ95] ———, *The lower algebraic K-theory of virtually infinite cyclic groups*, K-Theory **9** (1995), no. 1, 13–30. MR 1340838 (96e:19003)
- [Ji04] Lizhen Ji, *Asymptotic dimension and the integral K-theoretic Novikov conjecture for arithmetic groups*, J. Differential Geom. **68** (2004), no. 3, 535–544. MR 2144540 (2006c:57025)

- [Ji07] ———, *Integral Novikov conjectures and arithmetic groups containing torsion elements*, Comm. Anal. Geom. **15** (2007), no. 3, 509–533. MR 2379803 (2009b:22010)
- [Kas88] G. G. Kasparov, *Equivariant KK -theory and the Novikov conjecture*, Invent. Math. **91** (1988), no. 1, 147–201. MR 918241 (88j:58123)
- [Kas15] Daniel Kasprowski, *On the k -theory of groups with finite decomposition complexity*, Proceedings of the London Mathematical Society **110** (2015), no. 3, 565–592.
- [KLR] Holger Kammeyer, Wolfgang Lück, and Henrik Rüping, *The Farrell–Jones conjecture for arbitrary lattices in virtually connected Lie groups*, arXiv:1401.0876v1.
- [LR05] Wolfgang Lück and Holger Reich, *The Baum–Connes and the Farrell–Jones conjectures in K - and L -theory*, Handbook of K -theory. Vol. 1, 2, Springer, Berlin, 2005, pp. 703–842. MR 2181833 (2006k:19012)
- [Lüc00] Wolfgang Lück, *The type of the classifying space for a family of subgroups*, J. Pure Appl. Algebra **149** (2000), no. 2, 177–203. MR 1757730 (2001i:55018)
- [Lüc05] ———, *Survey on classifying spaces for families of subgroups*, Infinite groups: geometric, combinatorial and dynamical aspects, Progr. Math., vol. 248, Birkhäuser, Basel, 2005, pp. 269–322. MR 2195456 (2006m:55036)
- [Möh87] Walter Möhres, *Torsion-free nilpotent groups with bounded ranks of the abelian subgroups*, Group theory (Bressanone, 1986), Lecture Notes in Math., vol. 1281, Springer, Berlin, 1987, pp. 115–117. MR 921696
- [PW85] Erik K. Pedersen and Charles A. Weibel, *A nonconnective delooping of algebraic K -theory*, Algebraic and geometric topology (New Brunswick, N.J., 1983), Lecture Notes in Math., vol. 1126, Springer, Berlin, 1985, pp. 166–181. MR 802790 (87b:18012)
- [STY02] G. Skandalis, J. L. Tu, and G. Yu, *The coarse Baum–Connes conjecture and groupoids*, Topology **41** (2002), no. 4, 807–834. MR 1905840 (2003c:58020)
- [War76] Robert B. Warfield, Jr., *Nilpotent groups*, Lecture Notes in Mathematics, Vol. 513, Springer-Verlag, Berlin-New York, 1976. MR 0409661 (53 #13413)
- [Weg12] Christian Wegner, *The K -theoretic Farrell–Jones conjecture for $CAT(0)$ -groups*, Proc. Amer. Math. Soc. **140** (2012), no. 3, 779–793. MR 2869063
- [Weg15] ———, *The Farrell–Jones conjecture for virtually solvable groups*, J. Topol. **8** (2015), no. 4, 975–1016. MR 3431666

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY
E-mail address: kasprowski@mpim-bonn.mpg.de